

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 023 837 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
02.08.2000 Bulletin 2000/31

(21) Application number: **00300666.5**

(22) Date of filing: **28.01.2000**

(51) Int. Cl.⁷: **A01N 43/50, A01N 43/653**
// (A01N43/50, 43:90, 35:04,
25:30, 25:02), (A01N43/653,
43:90, 35:04, 25:30, 25:02)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **29.01.1999 US 240634**

(71) Applicant:
American Cyanamid Company
Madison, New Jersey 07940-0874 (US)


(72) Inventor: **Aven, Michael**
55131 Mainz (DE)

(74) Representative:
Walters, Philip Bernard William et al
Wyeth Laboratories,
Patents & Trade Marks Department,
Huntercombe Lane South,
Taplow
Maidenhead, Berkshire SL6 0PH (GB)

(54) **Non-aqueous, emulsifiable concentrate formulation**

(57) The invention relates to a non-aqueous, emulsifiable concentrate (EC) formulation for crop protection active compounds which comprises

(a1) 50 to 300 g/L of at least one azole derivative having a free hydroxy group or a salt or an adduct thereof, preferably a compound of formula I,



wherein, R¹ and R² each independently represent hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl or alkadienyl group;

R³ represents a halogen atom or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, alkoxy or aryl group;

A represents a nitrogen atom or a CH group; and

n represents an integer from 0 to 2;

(a2) optionally, 50 to 500 g/L of at least one additional fungicidally active compound;

(b) up to 700 g/L of one or more alkoxylates of an aliphatic alcohol,

(c) up to 100 g/L of one or more non-ionic dispersants,

(d) 10 to 100 g/L of one or more anionic dispersants,

(e) 50 to 600 g/L of one or more polar aprotic organic solvents, and

(f) 150 to 500 g/L of one or more non-polar organic solvents, any

EP 1 023 837 A2

(g) up to 5 g/L of one or more defoamers,

and to the use of such a emulsifiable concentrate as a fungicide.

Description

BACKGROUND OF THE INVENTION

[0001] The present invention relates to a non-aqueous, emulsifiable concentrate (EC) formulation for fungicidal azole compounds which comprises one or more fungicidal crop protection active compounds, one or more alkoxylates of an aliphatic alcohol, optionally one or more non-ionic dispersants, one or more anionic dispersants, one or more polar aprotic organic solvents, one or more non-polar organic solvents, and optionally one or more defoamers.

[0002] Emulsifiable concentrate (EC) formulations conventionally contain an active ingredient, one or more surfactants which act as emulsifiers upon dilution with water and a water immiscible solvent. Typical solvents for conventional EC formulations are aromatic hydrocarbons as for example xylene, Shellsol A or Solvesso 200. These solvents have very low solubilities in water and a high capability of dissolving a wide range of active ingredients.

[0003] Due to the presence of the solvent, many fungicides formulated as EC formulations have advantages such a higher degree of systemicity and overall activity compared to the same fungicide formulated as a wettable powder (WP), water dispersible granule (WG) or suspension concentrate (SC).

[0004] The observed efficacy of the combination of ingredients can sometimes be significantly higher than that would be expected from the amounts of the individual ingredients used (synergism). The efficacy of the active components can often be improved by addition of other ingredients such as adjuvants.


[0005] In order to increase the ease and safety of handling and dosing of these adjuvants by the end-user, and avoid unnecessary packaging material, it is desirable to develop concentrated formulations which already contain such adjuvants.

[0006] It is known that the activity of metconazole can be enhanced with certain adjuvants, in particular, with alkoxylated alcohols, as shown, for example, by U.S. Patent No. 5,393,770.

[0007] However, there is no suggestion of concentrated EC formulations comprising metconazole and alkoxylated alcohols.

SUMMARY OF THE INVENTION

[0008] Surprisingly, it has been found that stable EC formulations for fungicidal azole compounds having a free hydroxy group, such as bitertanol, cyproconazole, diniconazole, flutriazole, hexaconazole, tebuconazole, triadimenol, triticonazole and uniconazole, and preferably a compound of formula I



in which

R¹ and R² each independently represent hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl or alkadienyl group;

R³ represents a halogen atom or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, alkoxy or aryl group;

A represents a nitrogen atom or a CH group; and

n represents an integer from 0 to 2;
or a salt or an adduct thereof;

can be prepared if the formulation comprises, in addition to the compound of formula I, one or more alkoxylates of an aliphatic alcohol, one or more non-polar organic solvents, and at least one polar aprotic organic solvent.

[0009] The present invention therefore includes a non-aqueous, emulsifiable concentrate (EC) formulation for fungicidal crop protection active compounds which comprises

(a)

- (a1) 50 to 300 g/L of at least one azole derivative with a free hydroxy group or a salt or an adduct thereof;
 (a2) optionally 50 to 500 g/L of at least one additional fungicidally active compound

- (b) 100 to 700 g/L of one or more alkoxylate of an aliphatic alcohol,
 (c) up to 100 g/L of one or more non-ionic dispersants,
 (d) 10 to 100 g/L of one or more anionic dispersants,
 (e) 50 to 600 g of one or more polar aprotic organic solvents, and
 (f) up to 500 g/L of one or more non-polar organic solvents; and
 (g) up to 5 g/L of one or more defoamers.

[0010] The present invention also includes a method for combating a fungus at a locus which comprises emulsifying a formulation according to the present invention with water and treating said locus with the obtained diluted aqueous formulation.


DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0011] Preferred fungicidal crop protection compounds are those azoles of formula I, in which

- R^1 and R^2 each independently represent hydrogen atom or an alkyl group;
 R^3 represents a halogen atom or an optionally substituted alkyl group;
 A represents a nitrogen atom; and
 n represents 1; or a salt or an adduct thereof.

[0012] Particularly preferred are those compounds of formula I wherein A represents a nitrogen atom; R^1 and R^2 represent a C_{1-6} alkyl group, preferably a methyl group; and R^3 is attached in the para-position and represents a fluoro or chloro atom or a C_{1-6} haloalkyl group.

[0013] Most preferred is metconazole, a compound of formula IA,



which is known from "The Pesticide Manual," 10th Edition, The British Crop Protection Council and The Royal Society of Chemistry, 1994, (hereinbelow abbreviated as "Pesticide Manual"), page 669.

[0014] The compound of formula I, due to the basic nature of the azole ring, is capable of forming salts or addition products with inorganic or organic acids or metal ions. The compounds of formula I are preferably used as such in the EC formulation according to the present invention.


[0015] The compositions of this invention can be applied to plants or their environment simultaneously with or in succession with other active substances. These other active substances can be either fertilizers, agents which donate trace elements or other preparations which influence plant growth. However, they can also be herbicides, insecticides, fungicides, bactericides, nematocides, algicides, molluscicides, rodenticides, virucides, compounds inducing resistance into plants, biological control agents such as viruses, bacteria, nematodes, fungi and other microorganisms, repellents of birds and animals, and plant growth regulators, or mixtures of several of these preparations, if appropriate, together with other substances conventionally used in the art of formulation, such as surfactants or other additives which promote formulation stability.

[0016] The other fungicidal compound can be, for example, one which is capable of combating diseases of cereals (e.g. wheat) such as those caused by *Erysiphe*, *Puccinia*, *Septoria*, *Gibberella*, *Fusarium* and *Helminthosporium* spp., seed and soil borne diseases and downy and powdery mildews on vines and powdery mildew and scab on apples etc.

These mixtures of fungicides can have a broader spectrum of activity than the compound of general formula I alone.

[0017] Examples of the other fungicidal compounds are AC 382042, anilazine, azoxystrobin, benalaxyl, benomyl, binapacryl, blasticidin S, Bordeaux mixture, bromuconazole, bupirimate, captafol, captan, carbendazim, carboxin, carpropamid, chlorbenzthiazon, chlorothalonil, chlozolinate, cycloheximide, cymoxanil, cyprofuram, cyprodinil, dichlofluanid, dichlone, dichloran, diclobutrazol, diclocymet, diclomezine, diethofencarb, difenoconazole, diflumetorim, dimethirimol, dimethomorph, dinocap, ditalimfos, dithianon, dodemorph, dodine, edifenphos, epoxiconazole, etaconazole, ethirimol, etridiazole, famoxadone, fenapanil, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenpiclonil, fenpropidin, fenpropimorph, fentin, fentin acetate, fentin hydroxide, ferimzone, fluazinam, fludioxonil, flumetover, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, fuberidazole, furalaxyl, furametpyr, guazatine, IKF-916, imazalil, iminoctadine, ipconazole, iprodione, isoprothiolane, iprovalicarb, kasugamycin, KH-7281, kitazin P, kresoxim-methyl, mepanipyrin, mepronil, metalaxyl, methfuroxam, MON 65500, myclobutanil, neoasozin, nickel dimethyldithiocarbamate, nitrothalisopropyl, nuarimol, ofurace, organo mercury compounds, oxadixyl, oxycarboxin, penconazole, pencycuron, phenazineoxide, phthalide, polyoxin D, polyram, probenazole, prochloraz, procymidione, propamocarb, propiconazole, propineb, pyrazophos, pyrifenoxy, pyrimethanil, pyroquilon, pyroxyfur, quinomethionate, quinoxifen, quintozone, spiroxamine, SSF-126, SSF-129, streptomycin, sulfur, tecloftalame, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tolclofsmethyl, tolylfluanid, triadimefon, triazbutil, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, validamycin A, vinclozolin, XRD-563 and zarilamid.

[0018] In a preferred embodiment of the present invention the EC comprises a mixture of at least one compound of formula I and at least one compound of formula II,



in which

R^4 and R^5 each independently represent hydrogen or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, haloalkyl, aryl, heteroaryl, cycloalkyl, bicycloalkyl or heterocyclyl group, or

R^4 and R^5 together with the interjacent nitrogen atom represent an optionally substituted heterocyclic ring,

R^6 represents a halogen atom or an alkyl or alkoxy group,

m represents an integer from 0 to 5, and

Hal represents a halogen atom.

[0019] The compounds of formula II are known, for example, from U.S. Patent No. 5,593,996.

[0020] Preferred are those compounds of formula II, in which

R^4 represents an C_{1-8} alkyl, C_{2-8} alkenyl, C_{3-8} cycloalkyl or C_{1-8} haloalkyl group, and R^5 represents a hydrogen atom or a C_{1-8} alkyl group; or

R^4 and R^5 together with the interjacent nitrogen atom represent a C_{5-7} heterocyclic ring being optionally substituted by one or two C_{1-4} alkyl groups,


R^6 independently represent a fluorine or chlorine atom or a methoxy group,

m represents an integer from 2 or 3, and

Hal represents a chlorine atom.

[0021] Most preferred is 5-chloro-6-(2,4,6-trifluorophenyl)-7-[2-(1,1,1-trifluoro)propylamino]-[1,2,4]triazolo[1,5-a]pyrimidine, coded as "azolopyrimidine IIA" hereinbelow.

[0022] In another preferred embodiment of the present invention the EC comprises a mixture of at least one compound of formula I and at least one compound of formula III,



wherein

- 15 R^7 represents a halogen atom, an optionally substituted alkyl, alkanoyloxy or alkoxy group; or a hydroxy group,
 R^8 represents a halogen atom or an optionally substituted alkyl group,
 n is 0 or an integer of 1 to 3;
 R^9 independently represents a halogen atom, an optionally substituted alkyl or alkoxy group or a nitro group;
 R^{10} represents a halogen atom, a cyano, carboxy, hydroxy or nitro group or an optionally substituted alkyl, alkoxy,
 20 alkenyl, alkylthio, alkylsulphinyl, alkylsulphonyl or amino group;
 R^{11} represents an optionally substituted alkyl group;
 R^{12} represents a halogen atom or a nitro group, an optionally substituted alkyl, alkoxy, alkenyloxy, alkynyloxy,
 alkylthio, cycloalkyl, cycloalkyloxy, aryloxy group;
 r is 0, 1 or 2; and
 25 R^{13} independently represents a halogen atom, an optionally substituted alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy,
 alkynyloxy, cycloalkyl, cycloalkoxy group.

[0023] Preferred are those compounds of formula III, in which

- 30 R^7 represents a halogen atom, an alkyl or alkoxy group;
 R^8 represents a halogen atom or an alkyl group,
 n is 1;
 R^9 is attached in the ortho-position with respect to R^8 and represents a halogen atom,
 R^{10} represents an alkyl group;
 35 R^{11} represents an alkyl group;
 R^{12} and R^{13} independently represent an alkoxy group or a benzyloxy group, in which the phenyl ring may be substituted by one or more halogen atoms or alkyl or alkoxy groups; and
 r represents 1; and R^{13} is attached to the ortho-position with respect to R^{12} .

40 **[0024]** Most preferred is 3-bromo-2,2'-dimethyl-4',5',6,6'-tetramethoxybenzophenone coded benzoylbenzene IIIA.

[0025] The compounds of formula III are known, for example, from U.S. Patent No. 5,679,866.

[0026] The compounds of formula I exhibit an extraordinary efficacy against a broad range of phytopathogenic fungi, in particular against fungi from the classes Ascomycetes, Basidiomycetes, Phycmycetes and Deuteromycetes. They are systemic and may be applied as a leaf or soil fungicide.

45 **[0027]** The formulation according to the invention may be preferably applied for controlling the following phytopathogenic fungal species of the genera: *Alternaria*, *Botrytis*, *Cercospora*, *Colletotrichum*, *Erysiphe* (*Blumeria*), *Elsinoe*, *Fusarium*, *Gibberella*, *Guignardia*, *Helminthosporium*, *Hemileia*, *Monilinia*, *Mycosphaerella*, *Nectria*, *Phythium*, *Phytophthora*, *Plasmopara*, *Podosphaera*, *Pseudocercospora*, *Pseudoperonospora*, *Puccinia*, *Pyrenophora*, *Pyricularia*, *Rhizoctonia*, *Sclerotinia*, *Sclerotium*, *Septoria*, *Sphaerotheca*, *Tilletia*, *Typhula*, *Uncinula*, *Uromyces*, *Ustilago*,
 50 *Venturia*, *Verticillium* and others.

[0028] The application rate of the compound of formula I according to this invention is usually in the range of 1 to 500 grams of active ingredient (g a.i.) per hectare (h), with rates between 15 to 200 g a.i./ha generally achieving satisfactory control. The optimal rate for a specific application will depend on the crop(s) under cultivation and the predominant species of infesting fungus, and readily may be determined by established biological tests known to those skilled
 55 in the art.

[0029] The alkoxyates of aliphatic alcohols (b) are preferably liquid, semi-solid, waxy or solid polyalkoxylated aliphatic alcohols. These adjuvants are, as a rule, obtainable by alkoxylation of fatty alcohols having 5 - 20, preferably 7 - 19, and, in particular 9 - 14, C-atoms with an alkyleneoxide having 2 - 6, preferably 2 - 3 C-atoms, in particular, with

a mixture of ethylenoxide and propyleneoxide. The aliphatic moieties of the said fatty alcohols and amines may be straight-chained or branched. Preferably these compounds correspond to mixed random or block oligomers of the following formula



in which

the average of the indexes given is as follows:

- n is an integer from 5 to 20, in particular, 7 to 19;
 x is an integer from 1 to 20, in particular, 4 to 10; and
 the sum of y and z is an integer from 0 to 12, in particular, 0 to 10.

[0030] Of particular interest are those polyalkoxylated aliphatic alcohols which are liquids at temperatures down to at least 20°C having a viscosity of 30 to 100, in particular 50 to 80 mPa · s at 25°C. The compounds which are commercially available under the trademark Synperonic® and certain Atplus®-types (both sold by Uniqema, formerly ICI Surfactants), in particular, Synperonic® 91-6 and Atplus® MBA 11-7, have been proven to be especially advantageous.

[0031] In preferred embodiment of the present invention the non-ionic dispersant (c) is triglyceride, a polyoxyalkylene fatty acid or a polyoxyalkylene alkyl phenol. These dispersants, are as a rule, obtainable by alkoxylation of triglycerides, fatty acids or phenols. The alkoxylation of triglycerides results in mixtures of compounds with one to three glyceride side chains having 9 - 24, preferably 12 - 22, and, in particular 14 - 20, C-atoms, in particular with ethyleneoxide. The aliphatic moieties of the said triglycerides may be straight-chained or branched. Preferably, these compounds correspond to mixed oligomers resulting from the alkoxylation of castor or canola oil.

[0032] Further preferred dispersants (c) are, for example, Arkopal®-type alkylarylethoxylates (sold by Clariant GmbH, formerly Hoechst AG).

[0033] Other particularly preferred dispersants (c) are a castor oil ethoxylates, for example, Ukanil® 2507, which is commercially available from Henkel KGaA.

[0034] Still further preferred dispersants (c) are polyethyleneoxide-polypropyleneoxide block-copolymers, as for example, Pluronic®-type block-copolymers, which are available from BASF AG.

[0035] The anionic dispersants (d) are, as a rule, an alkali or earth alkali sulfonate, which includes also highly concentrated mixtures of such an alkali or earth alkali sulfonates with an organic diluent such as an alcohol, preferably, butanol or 2-ethylhexanol or aromatic hydrocarbons, preferably Solvesso® 200. Such a mixture preferably consists of 40 to 90 wt-% of at least one alkali or earth alkali sulfonate and 10 to 60 wt-% of an organic diluent. Ammonium, alkali and earth alkali alkylbenzene sulfonates are preferred, in particular, calcium dodecylbenzene sulfonates such as Rhodocal® 70/B (Rhodia, formerly Rhône-Poulenc), Phenylsulfonat CA100 (Clariant GmbH) or isopropylammonium dodecyl benzene sulfonate such as Atlox® 3300B (Uniqema).

[0036] Preferred polar aprotic solvents (e) can be compounds which are immiscible with water (e1) and have a dielectricity constant of at least 15 at 25 °C. Particularly preferred are N-alkylpyrrolidones such as N-octylpyrrolidone or alkyl lactates.

[0037] Another group of polar aprotic solvents (e) are compounds which are water-miscible (e2) and have a dielectricity constant of at least 15 at 25°C. Preferred are lactones such as γ-butyrolactone, ketones such as cyclohexanone, and N-cyclohexylpyrrolidone.

[0038] The solvent (f) is, as a rule, a water immiscible solvent in which the solubility of the crop protection compound (a) is greater than 5 g/L. Preferably (f) is a nonpolar organic solvent selected from the group consisting of aromatic hydrocarbons, aliphatic hydrocarbons, glycols and plant oil esters or mixtures thereof Preferred aromatic hydrocarbons are, e.g., toluene, xylenes, or substituted naphthalenes, as for example, solventnaphtha, Solvesso® 200 (Deutsche Exxon Chemicals) or Shellsol® A (Deutsche Shell AG). Preferred aliphatic hydrocarbons are, e.g. cyclohexane, paraffins as, for example, Isopar® H (Deutsche Exxon Chemicals) or Shellsol® T (Deutsche Shell AG), preferred plant oil esters are methylated coconut or soybean oil esters, in particular, methyl caprylate such as Witconol 1095 (Witco Corp.), preferred glycols are monoalkyl and dialkyl dialkeneglycols, in particular dimethyl diethyleneglycol (Diglyme), diethyl diethyleneglycol (Ethyl Diglyme) and monopropyl dipropyleneglycol such as Dowanol® DPNP Glycol Ether (Dow Chemical Company Ltd.). Mixtures of different liquids are often suitable.

[0039] Preferred anti-foam agents (g) are silica and polydialkylsiloxanes, in particular, polydimethylsiloxanes or mixtures thereof, such as Rhodorsil® 416 or Rhodosil® 454 from Rhodia, fluoroaliphatic esters such as Fluorad® FC-430 from 3M or perfluoroalkylphosphonic / perfluoroalkylphosphonic acids or the salts thereof such as Fluowet® PL80, or Fluowet® PP from Clariant. Particularly preferred is a combination of polydimethylsiloxanes and perfluoroalkylphosphonic / perfluoroalkylphosphonic acids.

[0040] Preferred are EC formulations comprising:

- (a1) 50 to 300 g/L of metconazole
- (a2) optionally, 50 to 300 g/L of at least one additional fungicidally active compound selected from the formulae II and III;
- (b) 100 to 700 g/L of one or more alkoxyate of an aliphatic alcohol, preferably, a C₅₋₂₀ alcohol alkoxyated with one to nine C₂₋₆ alkoxy groups;
- (c) up to 100 g/L of a non-ionic dispersant, preferably, a polyoxyethylene fatty acid,
- (d) 10 to 100 g/L of an anionic dispersant, in particular an amino sulfonate or an alkali or earth alkali sulfonate,
- (e) 50 to 600 g/L of one or more polar aprotic organic solvents, preferably, selected from the group consisting of N-C₂₋₁₆ alkylpyrrolidones, N-cycloalkylpyrrolidines, N-hydroxyalkylpyrrolidones and lactones; and
- (f) 160 to 500 g/L of one or more non-polar organic solvents, preferably, selected from the group consisting of diethyleneglycol dialkylethers, aromatic hydrocarbons and aliphatic hydrocarbons or mixtures thereof;
- (g) up to 5 g/L of a defoamer; preferably, a perfluoroalkyl phosphonic acid, a perfluoroalkyl phosphinic acid or a mixture thereof.

[0041] In a particularly preferred embodiment of this invention the EC formulation consists essentially of

- 50 to 250 g/L, preferably 20 to 100 g/L of an azole derivative of formula I, most preferably, metconazole;
- up to 200 g/L, preferably 20 to 150 g/L of a second fungicidal compound agent, most preferably, a compound of formula II or III;
- 150 to 500 g/L, preferably 200 to 450 g/L of one or two alkoxyates of an aliphatic alcohol, most preferably, a C₇₋₁₉ alcohol being alkoxyated with 4 to 10 ethoxy groups;
- 0 to 75 g/L, preferably 0 to 50 g/L of an ethoxylated fatty acid or phenol, most preferably, castor oil ethoxylate or nonylphenol ethoxylate;
- 5 to 100 g/L, preferably 10 to 75 g/L of an alkylbenzene sulfonate, most preferably, ammonium, potassium or sodium dodecylbenzene sulfonate;
- 100 to 500 g/L, preferably 120 to 480 g/L of a n-C₂₋₁₂ alkylpyrrolidone, most preferably, n-octylpyrrolidone, or γ -butyrolactone;
- 150 to 500 g/L, preferably 200 to 450 g/L of a non-polar organic solvent, most preferably, selected from the group consisting of diethyleneglycol dialkylethers, aromatic hydrocarbons and aliphatic hydrocarbons or mixtures thereof;
- up to 5 g/L, preferably 0.5 to 2 g/L of a defoamer selected from perfluoroalkyl phosphonic acids, perfluoroalkyl phosphinic acids and mixtures thereof;
- up to 2 g/L of a silicone-based defoamer.

[0042] A method of making such a composition is also provided which comprises bringing a compound of formula I into association with the ingredients (b) to (g). It is also envisaged that different isomers or mixtures of isomers of formula I may have different levels or spectra of activity and thus compositions may comprise individual isomers or mixtures of isomers.

[0043] Pesticidal compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application.

[0044] The compositions of the invention may contain 0 to 10% w/v of additives besides (b) to (f) such as corrosion inhibitors, stabilizers, penetrants and stickers. Certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation and crystallization.

[0045] Aqueous emulsions, for example, compositions obtained by diluting the EC formulated product according to the invention with water, also lie within the scope of the invention.

[0046] As a commodity, the compositions are in a concentrated form, whereas the end user generally employs diluted compositions. The compositions may be diluted to a concentration down to 0.001% of active ingredient. The doses usually are in the range from 0.001 to 10 kg a.i./ha, preferably 0.03 to 0.5 kg a.i./ha, and most preferably, 0.04 to 0.4 kg a.i./ha.

[0047] For a clearer understanding of the invention, specific examples are set forth below. These examples are merely illustrations and are not to be understood as limiting the scope and underlying principles of the invention in any way. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the following examples and foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

EXAMPLES:

[0048] Examples of formulations according to the invention are shown in the following examples A to M:

5 Identity of Ingredients used in Examples

[0049]

10

15

20

25

Name	Identity
Metconazole	Fungicidal azole of formula IA
Azolopyrimidine IIA	Fungicidal triazolopyrimidine of formula II
Benzoylbenzene IIIA	Fungicidal benzoylbenzene of formula III
Synperonic® 91-6 (Uniqema)	Alcohol ethoxylate
Synperonic® NP-4 (Uniqema)	Nonylphenol ethoxylate
Atplus MBA 11-7® (Uniqema)	Monobranched Alcohol ethoxylate
Rhodocal® 70/B (Rhodia)	70% Calcium Dodecylbenzene sulfonate in butanol
Atlox® 3300B (Uniqema)	Isopropylammonium Dodecylbenzene sulfonate
Ukanil® 2507 (Uniqema)	Castor oil ethoxylate
Fluowet® PL80 (Clariant GmbH)	Mixture of perfluoroalkylphosphonic and perfluoroalkylphosphinic acids
Mergital EL33 (Henkel)	Castor oil ethoxylate with 33 EO units

30 **Examples A and B**

[0050] All ingredients are weighed into a container and stirred until a homogenous solution is obtained.

35

40

45

Example A		Example B	
Ingredient	Concentration (g/L)	Ingredient	Concentration (g/L)
Metconazole	90	Metconazole	90
Atlox 3300B	50	Atlox 3300B	50
Ukanil 2507	20	Ukanil 2507	20
Synperonic 91-6	480	Synperonic 91-6	480
N-octylpyrrolidone	200	N-dodecylpyrrolidone	200
Solventnaphtha	to 1 liter	Solventnaphtha	to 1 liter

Physico-chemical Tests

50

[0051]

55

Phys-chem Tests	Example A	Example B
Density	0.97 g/ml	0.96 g/ml

(continued)

Phys-chem Tests	Example A	Example B
Flash point	59°C	60°C
Spray dilution 0.5 hours	ok (homogenous, no cream or precipitate)	ok
Spray dilution 2 hours	ok	ok
Spray dilution 4 hours	ok	ok
Spray dilution 24 hours	ok	ok
Storage of EC (7 days 0°C)	no crystals in EC	no crystals in EC
Storage of EC (14 days 40°C)	no crystals in EC, spray dilution as above ok	no crystals in EC, spray dilution as above ok
Storage of EC (14 days 54°C)	no crystals in EC, spray dilution as above ok	no crystals in EC, spray dilution as above ok

Example C

[0052] All ingredients are weighed into a container and stirred until a homogenous solution is obtained.

Ingredient	Concentration (g/L)
Metconazole	60
Azolopyrimidine IIA	100
Synperonic 91-6	350
Synperonic NP-4	50
Rhodocal 70/B	50
Fluowet PL80	1
n-Octylpyrrolidone	to 1 liter

Physico-chemical Tests

[0053]

Phys-chem Tests	Example C
Density	1.01 g/mL
Flash point	> 83°C
Spray dilution in 100 ml graduated cylinder, self-emulsification checked, followed by 30 inversions, foam judged (0 hour)	good self-emulsification, after inversions no foam
Spray dilution 0.5 hours	ok
Spray dilution 2 hours	ok
Spray dilution 4 hours	ok

[0054] The EC formulation of Example C combines an adjuvant (alcohol ethoxylate) necessary for good perform-

EP 1 023 837 A2

ance of metconazole with a polar aprotic solvent (n-octylpyrrolidone) necessary to dissolve 100 g/L of azolopyrimidine IIA in a water immiscible system. It is possible to raise the metconazole concentration above 60 g/l.

Examples D to H

[0055] All ingredients are weighed into a container and stirred until a homogenous solution is obtained.

Ingredient	Concentration (g/L)				
	Example				
	D	E	F	G	H
metconazole	125	125	125	83	83
Benzoylbenzene IIIA	90	90	90	60	60
γ -Butyrolactone	100	-	-	-	-
Synperonic 91-6	300	300	-	-	-
Atplus MBA 11-7	-	-	300	-	-
Atlox 3300B	-	50	50	-	-
Rhodocal 70/B	40	-	-	50	50
Ukanil 2507	50	30	30	30	30
Fluowet PL80	1	1	1	-	-
N-Cyclohexylpyrrolidone	-	-	-	240	240
N-Octylpyrrolidone	-	150	150	240	120
Solventnaphtha	100	to 1 liter	to 1 liter	to 1 liter	to 1 liter
Diethyleneglycol dimethylether	to 1 liter	-	-	-	-

[0056] Diethyleneglycol diethylether can also be used to partially replace diethylenglycol dimethylether to increase the flash point.

Physico-chemical tests**[0057]**

5

10

15

20

25

30

Phys-chem Tests	Example				
	D	E	F	G	H
Density (g/ml)	1.04	1.00	1.00	0,98	0,98
Flash point	55°C	54°C	60°C	54°C	52°C
Spray dilution 0.5 h	ok *	ok	ok	ok	ok
Spray dilution 1 h	ok	ok	ok	ok	ok
Spray dilution 2 h	ok	ok	ok	ok	ok
Spray dilution 4 h	ok	ok	ok	ok	ok
Spray dilution 20 h	not recorded	not recorded	not recorded	≈ 1 ml cream, otherwise ok	≈ 1 ml cream, otherwise ok
Spray dilution 24 h	some crystals as precipitate	ok, even after 4 days	very few crystals as precipitate	not tested	not tested
Storage of EC (7 days 0°C)	no crystals	crystals, clear	no crystals	no crystals	no crystals
Storage of EC (7 days -5°C)	no crystals	crystals	some crystals	no crystals	no crystals

* Ok means that the composition is homogenous, there is no cream or precipitate.

[0058] The EC formulations of Examples D to H combine an adjuvant (alcohol ethoxylate) necessary for good performance of metconazole with a polar aprotic solvent necessary to dissolve 125 g/l of benzoylbenzene IIIA in a water immiscible system. It is possible to raise the metconazole concentration above 90 g/l.

35

Example I

[0059] An EC formulation is prepared containing:

40

45

50

55

Ingredient	Concentration (g/L)
Metconazole	90
Benzoylbenzene IIIA	100
Synperonic 91-6	300
Phenylsulfonat CA100	30
Mergital EL33	60
Fluowet PL80	0.5
Rhodorsil 454	0.2
Solventnaphtha	175
γ-Butyrolactone	170
Solvesso 200	to 1 L


[0060] The new ECs are biologically very active (much more than ECs without an adjuvant). Most of the spray dilutions (emulsions) are stable despite a high concentration of water miscible substances (Synperonic 91-6, γ -butyrolactone, N-cyclohexylpyrrolidone). The ingredients have a good environmental profile. In the past, adjuvants have typically been added to the spray tank separately from the pesticidal formulation ("tank-mix adjuvant"). The adjuvant in a one-pack formulation is easier to use than as a tank-mix adjuvant.

Claims

1. A non-aqueous, emulsifiable concentrate (EC) formulation for fungicidal crop protection active compounds which comprises

- (a1) 50 to 300 g/L of at least one azole derivative having a free hydroxy group or a salt or an adduct thereof;
- (a2) optionally 50 to 500 g/L of at least one additional fungicidally active compound;
- (b) 100 to 700 g/L of one or more alkoxyates of an aliphatic alcohol,
- (c) up to 100 g/L of one or more non-ionic dispersants,
- (d) 10 to 100 g/L of one or more anionic dispersants,
- (e) 50 to 600 g/L of one or more polar aprotic organic solvents, and
- (f) up to 500 g/L of one or more non-polar organic solvents, and
- (g) up to 5 g/L of one or more defoamers.

2. A formulation according to Claim 1 wherein component (a1) is a compound of formula I




in which

- R^1 and R^2 each independently represent hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl or alkadienyl group;
- R^3 represents a halogen atom or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, alkoxy or aryl group;
- A represents a nitrogen atom or a CH group; and
- n represents an integer from 0 to 2;

3. A formulation according to Claim 1 wherein component (a1) is metconazole.


4. A formulation according to Claim 1 wherein the second active ingredient (a2) is a triazolopyrimidine of formula II



in which

R^4 and R^5 each independently represent hydrogen or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, haloalkyl, aryl, heteroaryl, cycloalkyl, bicycloalkyl or heterocyclyl group, or
 R^4 and R^5 together with the interjacent nitrogen atom represent an optionally substituted heterocyclic ring,
 R^6 represents a halogen atom or an alkyl or alkoxy group,
 m represents an integer from 0 to 5, and
Hal represents a halogen atom.

5. A formulation according to Claim 1 wherein the second active ingredient (a2) is a benzoylbenzene of formula III



wherein

R^7 represents a halogen atom, an optionally substituted alkyl, alkanoyloxy or alkoxy group; or a hydroxy group,
 R^8 represents a halogen atom or an optionally substituted alkyl group,
 n is 0 or an integer of 1 to 3;
 R^9 independently represents a halogen atom, an optionally substituted alkyl or alkoxy group or a nitro group;
 R^{10} represents a halogen atom, a cyano, carboxy, hydroxy or nitro group or an optionally substituted alkyl, alkoxy, alkenyl, alkylthio, alkylsulphinyl, alkylsulphonyl or amino group;
 R^{11} represents an optionally substituted alkyl group;
 R^{12} represents a halogen atom or a nitro group, an optionally substituted alkyl, alkoxy, alkenyloxy, alkynyloxy, alkylthio, cycloalkyl, cycloalkyloxy, aryloxy group;
 r is 0, 1 or 2; and
 R^{13} independently represents a halogen atom, an optionally substituted alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy, cycloalkyl, cycloalkoxy group.

6. A formulation according to Claim 1 wherein said alkoxylate of an aliphatic alcohol (b) is a C_{5-20} alcohol being alkoxylated with 1 to 20 C_{2-6} alkoxy groups.
7. A formulation according to Claim 5 wherein said the alkoxylate (b) is a straight-chained or branched C_{7-19} alcohol being ethoxylated with 4 to 18 ethoxy and/or propoxy groups, or a mixture thereof.
8. A formulation according to Claim 1 wherein the ratio of the crop protection active compounds (a) to said adjuvant (b) is between 1:0.5 and 1:100, preferably between 1:1 and 1:10.
9. A formulation according to Claim 1 wherein the non-ionic dispersant (c) is a polyoxyethylene fatty acid, or a polyoxyalkylene triglyceride derivative.
10. A formulation according to Claim 1 wherein the anionic dispersant (d) is an amino sulfonate or an alkali or earth alkali sulfonate.
11. A formulation according to Claim 1 wherein the polar aprotic solvent (e) is immiscible with water.
12. A formulation according to Claim 11 wherein the polar aprotic solvent (e) is selected from the group consisting of n - C_{2-16} alkylpyrrolidones, n -cycloalkylpyrrolidines, n -hydroxyalkyl-pyrrolidones and lactones.
13. A formulation according to Claim 1 wherein the non-polar solvent (f) is selected from the group consisting of diethyleneglycol dialkylethers, aromatic hydrocarbons and aliphatic hydrocarbons or mixtures thereof.
14. A formulation according to Claim 1 wherein the defomer (g) is selected from the group comprising perfluoroalkyl-

phosphonic acids, perfluoroalkylphosphinic acids and mixtures thereof.

15. An EC according to Claim 14 which additionally comprises a silicone-based defoamer.

5 16. A method for combating a fungus at a locus which comprises emulsifying a formulation as claimed in Claim 1 with water and treating said locus with the obtained diluted aqueous formulation.

10

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 023 837 A3

(12)

EUROPEAN PATENT APPLICATION(88) Date of publication A3:
30.05.2001 Bulletin 2001/22(43) Date of publication A2:
02.08.2000 Bulletin 2000/31

(21) Application number: 00300666.5

(22) Date of filing: 28.01.2000


(51) Int. Cl.⁷: **A01N 43/50**, A01N 43/653,
A01N 25/02// (A01N43/50, 43:90, 35:04,
25:30, 25:02), (A01N43/653,
43:90, 35:04, 25:30, 25:02)(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 29.01.1999 US 240634

(71) Applicant:
**American Cyanamid Company
Madison, New Jersey 07940-0874 (US)**(72) Inventor: **Aven, Michael
55131 Mainz (DE)**(74) Representative:
**Langfinger, Klaus Dieter, Dr. et al
Director Patents & Licensing,
BASF Aktiengesellschaft ZDX/P-C6
67056 Ludwigshafen (DE)**(54) **Non-aqueous, emulsifiable concentrate formulation**

(57) The invention relates to a non-aqueous, emulsifiable concentrate (EC) formulation for crop protection active compounds which comprises

(a1) 50 to 300 g/L of at least one azole derivative having a free hydroxy group or a salt or an adduct thereof, preferably a compound of formula I,

wherein, R¹ and R² each independently represent hydrogen atom or an optionally substituted alkyl, alkenyl, alkynyl or alkadienyl group;R³ represents a halogen atom or an optionally substituted alkyl, alkenyl, alkynyl, alkadienyl, alkoxy or aryl group;

A represents a nitrogen atom or a CH group; and

n represents an integer from 0 to 2;

(a2) optionally, 50 to 500 g/L of at least one additional fungicidally active compound;

(b) up to 700 g/L of one or more alkoxylates of an aliphatic alcohol,

(c) up to 100 g/L of one or more non-ionic dispersants,

EP 1 023 837 A3

EP 1 023 837 A3

- (d) 10 to 100 g/L of one or more anionic dispersants,
- (e) 50 to 600 g/L of one or more polar aprotic organic solvents, and
- (f) 150 to 500 g/L of one or more non-polar organic solvents, any
- (g) up to 5 g/L of one or more defoamers,

and to the use of such a emulsifiable concentrate as a fungicide.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 30 0666

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
P, X	EP 0 933 025 A (AMERICAN CYANAMID CO) 4 August 1999 (1999-08-04) * the whole document *	1-16	A01N43/50 A01N43/653 A01N25/02
X	EP 0 351 195 A (CHEMICCA LTD) 17 January 1990 (1990-01-17) * the whole document * * examples *	1-16	/(A01N43/50, 43:90, 35:04, 25:30, 25:02), (A01N43/653, 43:90, 35:04, 25:30, 25:02)
A	US 5 393 770 A (GRAYSON BASIL T) 28 February 1995 (1995-02-28)		
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A01N
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 4 April 2001	Examiner Bertrand, F
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 0666

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

04-04-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0933025 A	04-08-1999	BR 9900060 A	21-03-2000
		JP 11269006 A	05-10-1999
EP 0351195 A	17-01-1990	NZ 225428 A	26-03-1991
		AT 111300 T	15-09-1994
		AU 633497 B	04-02-1993
		AU 3793989 A	18-01-1990
		BR 8903512 A	13-03-1990
		DE 68918187 D	20-10-1994
		DE 68918187 T	03-08-1995
		ES 2059761 T	16-11-1994
		FI 893441 A	16-01-1990
		JP 2059501 A	28-02-1990
		PT 91183 A, B	08-02-1990
		ZA 8905274 A	25-04-1990
US 5393770 A	28-02-1995	EP 0526206 A	03-02-1993
		JP 7179306 A	18-07-1995

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82